

Soluble powder *

20% Sodium Sulfachloropyrazine Soluble Powder



Category

Soluble powder

Appearance

light yellow powder.

Application


same as functional description.

Good Manufacturing Practice



Functions

Used for the treatment of coccidiosis in livestock, poultry and rabbits, also used for the treatment of avian cholera and typhoid fever.



! Dosage and administration method

according to this product. Mixed Drinking: 1.5 grams per 1 liter of water for broilers and turkeys for 3 consecutive days. Mixed feeding: per 1000 kg of feed, 3000 g for broilers and turkeys for 3 consecutive days; 3000 g for rabbits for 3 consecutive days. Oral: 1.2 ml (formulated as a 10% solution) per 1 kg of body weight for sheep for 3 to 5 days.

🔔 Specification

20%.

Safety And Packaging

! Side effects

No adverse reactions at recommended doses.

🔔 Precautions

Mixed drinking not more than 5 days. Prohibit use during the laying period of hens. Prohibit mixing with feed for a long time.

! Dormancy period

sheep 28 days (tentative), turkey 4 days, broiler 1 day.



Storage

Shielded from light, airtight, kept in a dry place

Packing

100g/bag

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com