

Injection



10% Vitamin C Injection



Function

Nutritional medicine

Appearance

colorless to light yellow clarified liquid.

Good Manufacturing Practice



Indications

Used for vitamin C deficiency and various infectious diseases, high fever, trauma or burns, also used for anemia, hemorrhagic tendency, methemoglobinemia, atopic dermatitis auxiliary treatment.

Usage and dosage

Intramuscularly or intravenously, one-time dose of
Horse: 10 - 30 ml.
Cattle: 20 - 40 ml.
Sheep and pigs: 2 - 5 ml.
Dogs: 0.2 - 1 ml.

Specification

1.5 ml: 0.5 grams
2.10ml: 1 gram

Safety And Packaging

Side effects

Increased risk of uric acid, oxalic acid or cysteine crystal formation when used in high doses.

Precautions

1. Should not be mixed with alkaline drugs (e.g. sodium bicarbonate, etc.), vitamin B12, vitamin K3, as it is prone to oxidative failure.
2. Inactivates aminoglycosides, β -lactam antibiotics, tetracyclines, etc. to varying degrees. Therefore, it should not be mixed with these antibiotics for injection.
3. When used in large doses, it can acidify the urine and increase the excretion of certain organic base drugs.
4. Large doses can cause diarrhea and urinary calculi.



Dormant period

not specified.

Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com