

Tablets

*

Oxytetracycline Tablets

For the treatment of infectious diseases caused by sensitive gram-positive bacteria, negative bacteria, mycoplasma and so on.




Function

Antibacterial drug.

Appearance

light yellow tablet.

Good Manufacturing Practice



Indications

for the treatment of infectious diseases caused by sensitive gram-positive bacteria, negative bacteria, mycoplasma and other infectious diseases.

Dosage and Administration

Internal, one dose.
Pigs, foals, calves, lambs: 10mg-25mg/kg body weight.
Dogs: 15mg-50mg/kg body weight.
Poultry: 25mg-50mg/kg body weight.
Use 2-3 times daily for 3-5 days.

Specification

C22H24N2O9
50mg、12.5mg、250mg

Safety And Packaging

Side Effects

Localized irritation.
Disturbance of intestinal flora.
Impact on dental and bone development.
Liver and kidney damage.
Cardiovascular effects.
Anti-metabolic effects.

Precautions

Adult ruminants, horses and rabbits should not be used internally. Prolonged use may induce secondary infections. It is prohibited for animals with severe liver and kidney dysfunction. Avoid taking with dairy products and feeds with high calcium content.

Stopping period

Calf, sheep, pig: 7 days
Laying hens: 2 days
Milking cow: 72 hours

Storage

Sealed in dry place and keep away from light.

Packing

5 tablets/plate
Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com