

Injection



Amoxicillin 15% Injection



Function

antibiotic drug

Appearance

white or off-white oily suspension

Good Manufacturing Practice



Indications

respiratorytract infections

- Used for the treatment of respiratory tract infections caused by sensitive bacteria. such as bacterial pneumonia in pigs.

Digestive Tract Infections

- For the treatment of digestive tract infections such as bacterial enteritis
- Can be used to treat digestive tract infections in pigs. such as E. coli disease.

Genitourinary Tract Infections

- Used for the treatment of genitourinary tract infections such as uteritis and cystitis.
- Can be used to treat urinarytract infections in sows.

Skin and Soft Tissue Infections

- For the treatment of skin and soft tissue infections such as wound infections. abscesses. hoof infectionsIt can be used to treat streptococcal arthritis and soft tissue infections in pigs

Dosage

intramuscular and subcutaneous: 0.1 ml per kg of body weight in cattle. pigs, dogs and cats. and can be repeated after 48 hours if necessary.

Specification

15%

Safety And Packaging

Precautions

1. Shake well before use.
2. Do not drink milk during the treatment period and 96 hours before stopping.

Storage

Protected from light. sealed and kept in a cool place



Discontinuation period

28 days for cows and pigs. 96 days for milk abandonment

Packing

100ml1 bottle; 50ml/ bottle.

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com