

Injection



1% Ivermectin Injection



Function

Antiparasitic


Composition

10 mg of ivermectin per ml.

Appearance

colorless or almost colorless,
liquid, slightly viscous.

Good Manufacturing Practice



Usage and dosage

Subcutaneous injection;
Per kg of body weight: cattle, sheep: 0.02 ml;
pigs: 0.03 ml.

Specification

1%

Safety And Packaging

Side effects

- 24 hours after use, horses may experience allergic reactions caused by dead Onchocerca volvulus. Such as edema and itching around the injection site.
- Serious adverse reactions may occur when used in bullous fly disease (because the larvae are located in critical areas of the body).
- Shock associated with dead microfilariae may occur in dogs when killing microfilariae.
- Poultry may experience death, lethargy or loss of appetite.
- When injected, discomfort or temporary edema at the injection site may occur.

Precautions

- Cattle and sheep should not be treated within 21 days prior to slaughter; camels should not be treated within 28 days prior to slaughter.
- This product should not be used for intravenous or intramuscular injection.



Storage

Not specified in the current standard.

Packing

50 bottles/100 bottles/120 bottles/150 bottles/carton.

Packaging specifications can be adjusted
according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com