

Injection \*

## 10% Florfenicol Injection



### Function

Antibacterial

### Ingredients

100 mg of florfenicol per ml.

### Appearance

Colorless or almost colorless, liquid.  
slightly viscous.

## Good Manufacturing Practice



### Respiratory Diseases

- Infectious pleuropneumonia, wheezing disease, parvovirus in pigs.
- Respiratory infections of birds, such as cholera, white diarrhea, respiratory disease caused by Escherichia col, Salmonella, Bartonella.

### Digestive Diseases

- Piglets yellow diarrhea, white diarrhea, edema disease, enteritis.
- Poultry cholera, white diarrhea, diarrhea, intractable diarrhea.

### Sustemic Infections

- Streptococcal disease, staphylococcal disease, exudative dermatitis in pigs.
- Postpartum infection in pigs, such as endometritis, lactation disorder syndrome.

### Other Infections

- Bacterial meningitis, urinary tract infection in pigs.
- Infectious plasmacytitis, pericardium and liver of poultry.



### Dosage

Pig, chicken: 0.2 ml per kg of body weight.

### Dosage interval

Once in 48 hours, 2 times in a row.

## Safety And Packaging

### Side effects

Prohibited to use in milk-laying period and egg-laying chickens.



### Precautions


- For livestock with renal insufficiency, the dosage should be reduced or the dosage interval should be extended.
- Prohibit the use during the vaccination period or when the immune function is seriously low.



### Rest period

Pig: 14 days

Chicken: 28 days



### Packing

100ml bottle.

Packaging specifications can be adjusted according to customer requirements.

## Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613  
E-mail: hebeijinfeng@gmail.com