

Injection



10% Tylosin Tartrate Injection



Function

Antimicrobial

Ingredients

Each ml contains: Tylosin Tartrate 100 mg.

Appearance


colorless or slightly yellow transparent liquid.

Good Manufacturing Practice



Indications

- Infections caused by microorganisms sensitive to tylosin, such as respiratory infections in cattle, sheep and swine, Duchenne dysentery in pigs, dysentery and arthritis caused by Mycoplasma bovis, mastitis and endometritis.
- Hypersensitivity to tylosin and cross-sensitization to macrolides.



Usage and dosage

1. Intramuscular or subcutaneous injection.
2. Cattle: 0.25 - 0.5 ml per 10 kg body weight per day for 3 - 5 days.
3. Calves, sheep, goats: 0.7 - 2.1 ml per 50 kg body weight per day for 3 - 5 days.
4. Pigs: 0.25 - 0.4 ml per 10 kg body weight every 12 hours for 3 days. Dogs: 0.2 - 1 ml.
5. Dogs, cats: 0.25 - 1 ml per 10 kg bw per day for 3 - 5 days.

Specification

10%

Safety And Packaging

Side effects

The most common adverse reaction is local irritation at the injection site.

Rest period

Meat animals: 8 days.
Dairy animals: 4 days.

Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com