

Injection



5% Flunixin Meglumine Injection



Function

Antipyretic, analgesic, and anti-inflammatory drug

Appearance


Colorless to slightly yellow transparent solution

Good Manufacturing Practice



Indications

For Horses
For the relief of inflammation and pain associated with musculoskeletal disorders.
For the relief of visceral pain associated with acute peritonitis.
For Cattle
For the control of fever associated with respiratory diseases and endotoxemia.
For the control of inflammation associated with endotoxemia.



Usage and Dosage

Calculated based on flunixin meglumine.
Intramuscular injection or intravenous injection: Single dose,
Horses, cattle, pigs: 2 mg per kilogram of body weight.
Administer 1 - 2 times per day. Do not use for more than 5 consecutive days.

Specification

100 mL: 5 g

Safety And Packaging

Precautions

1. Use with caution in animals with gastrointestinal ulcers, kidney disease, liver disease, or a history of bleeding.
2. Use with caution in the treatment of acute peritonitis; it may mask the loss of intestinal motility and cardiopulmonary symptoms caused by endotoxemia.
3. Use with caution in pregnant animals.
4. Arterial injection is prohibited, as it may cause central nervous system stimulation, ataxia, hyperpnea, and muscle weakness.
5. Horses may experience potential gastrointestinal intolerance, hypoproteinemia, and congenital diseases. Dogs may experience reduced gastrointestinal function.

Withdrawal period

Cattle, pigs: 28 days



Storage

Keep sealed and protect from light.

Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com