

Injection



## 20% Tylosin Tartrate Injection



### Function

Antimicrobial


### Composition

Each ml contains: Tylosin (as tartrate) 200 mg.

### Appearance


colorless or yellowish transparent liquid.

## Good Manufacturing Practice



### Indications

Infections caused by microorganisms susceptible to tylosin, e.g., respiratory infections in cattle, sheep and swine, Duchenne dysentery in swine, dysentery and arthritis caused by Mycoplasma, mastitis and endometritis.



### Usage and dosage

1. Intramuscular or subcutaneous injection.
2. Cattle: 0.5 - 1 ml per 10 kg body weight per day for 3 - 5 days. Calves, sheep, goats: 1.5 - 2 ml per 50 kg bw per day for 3 - 5 days. 2 times a day Cattle: 0.5 - 1 ml per 10 kg body weight per day for 3 - 5 days. Calves, Dogs, cats: 0.5 - 2 ml per 10 kg bw per day for 3 - 5 days.

### Specification

20%.

## Safety And Packaging

### Side effects

The most common adverse reaction is local irritation at the injection site.

### Rest period

Meat animals: 8 days.

Dairy animals: 4 days.



### Storage

Protected from light, sealed. It is recommended to keep it dry at room temperature.

### Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

### Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613  
E-mail: hebeijinfeng@gmail.com