

Injection *

20% Procaine Penicillin Injection



Function

Antimicrobial


Composition

Each ml contains 200 mg of procaine penicillin.

Appearance


This product is an oily solution containing fine particles.

Good Manufacturing Practice



Indications

It is mainly used for the treatment of chronic infections caused by penicillin-sensitive strains of bacteria, such as uterine pus accumulation, mastitis, fractures in cattle, and infections caused by actinomycosis and leptospirosis.



Usage and dosage

Intramuscular injection, one-time dose.
 Horses, cattle: 0.05 – 0.1 ml per kg body weight.
 Sheep, pig, foal, calf: 0.1 – 0.15 ml per kg body weight.
 Dogs, cats: 0.15 – 0.2 ml per kg body weight.
 Once a day for 2-3 days.

Specification

20%.

Safety And Packaging

Side effects

1. Mainly allergic reactions, which may occur in most domestic animals, but the incidence is low. Local reactions at the injection site are characterized by edema and pain, and systemic reactions are urticaria and rash, which can lead to shock or death in severe cases.
2. Attention should be paid to interactions and incompatibilities with other drugs to avoid affecting the efficacy.



Dormancy period

Meat animals:
 Cattle: 10 days
 Sheep: 9 days
 Swine: 7 days
 Milking animals: 48 hours

Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
 E-mail: hebeijinfeng@gmail.com