

Injection *

Lincomycin Hydrochloride and Spectinomycin



Function

Antimicrobial

Ingredients

Each ml contains: Lincomycin Hydrochloride 50 mg and Spectinomycin Hydrochloride 100 mg

Appearance


Colorless or slightly yellow transparent liquid

Good Manufacturing Practice



Indications

For Gram-positive. Gram-negative and Mycoplasma infections, for the treatment of chronic respiratory diseases in poultry. dysentery. infectious arthritis. pneumonia. erythema in swine and bacterial enteritis and pneumonia in calves.



Usage and dosage

Subcutaneous injection. one-time dose

Poultry

30 mg per kg body weight (lincomycin and macrolide combined)
Intramuscular injection, single dose. pigs. calves. weight (lincomycin and macrolide combined sheep: 15 malka bodu

Safety And Packaging

Precautions

1. Intravenous injection is prohibited. Intramuscular injection should be carried out slowly.
2. Antagonistic effect with tetracyclines
3. Shock associated with dead microfilariae may occur in dogs when

Dormancy period

Meat: 28 days. Milk: 7 days.

Packing

100 ml/ bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com