



30% Florfenicol Injection



Function

Antimicrobial

Composition

300 mg florfenicol per ml.

Appearance

colorless or yellowish transparent liquid.

Good Manufacturing Practice

Indications

Respiratory diseases

• it is used for the treatment of respiratory infections caused by Actinobacillus porcine pleuropneumoniae, Pasteurella multocida, Mycoplasma and other sensitive bacteria, such as infectious pleuropneumonia, wheezing disease, parvovirus disease, and so on.

Systemic infections

• it can be used to treat mixed infections such as hyperthermia syndrome, respiratory syndrome, streptococcal disease, toxoplasmosis, erythrocytosis, etc. caused by sensitive bacteria.

Gastrointestinal and genitourinary tract infections

• it is used for the treatment of enteritis, diarrhea, urinary tract infections caused by sensitive bacteria.

Other infections

· it can also be used to treat endometritis and pelvic inflammatory disease of domestic animals after childbirth.diarrhea, urinary tract infections caused by sensitive bacteria.

Usage and dosage

Pigs and chickens: 0.067 ml per kg body weight. Dosage interval: once in 48 hours, 2 times in a row.

Safety And Packaging

O Side effects

- 1. For livestock with renal insufficiency, the dose should be reduced or the dosage interval should be prolonged.
- 2. Prohibit use during the vaccination period or when the immune function is severely depressed.



煹 Rest period

Pig: 14 days.



Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613 E-mail: hebeijinfeng@gmail.com