

Soluble powder *

10% Sodium Sulfachloropyrazine Soluble Powder

Antibiotic drugs



Application

same as functional description.

Ingredients

Sulfachloropyrazine sodium

Appearance


light yellow powder.

Good Manufacturing Practice



Functions

Used for the treatment of coccidiosis in livestock, poultry and rabbits, also used for the treatment of avian cholera, typhoid fever and so on.



Dosage and method of administration

according to this product. Mixed Drinking: 3 grams per 1 liter of water for chickens and turkeys for 3 consecutive days. Mixed feeding: 6000 grams per 1000 kg of feed for chickens and turkeys for 3 consecutive days; 6000 grams for rabbits for 5 to 10 consecutive days. Oral: 1.2 ml (formulated as a 10% solution) per 1 kg of body weight for sheep for 3 to 5 consecutive days.

Safety And Packaging

Side effects

There are no adverse reactions to the use of this product according to the instructions.

Precautions

1. Drinking water administration should not exceed 5 consecutive days.
2. Prohibit the use of hens during egg-laying period.
3. Prohibit the use of long-term addition to feed.

Dormancy period

Chicken: 8 days.



Storage

Sealed and stored in a cool, light-proof place.

Packing

100g/bag

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com