

Injection



Oxytetracycline Hcl 20% Injection



Function

Antibacterial drug


Ingredients

Mainly used for the treatment of infections with Gram-positive cocci and Gram-negative bacilli sensitive to penicillin.

Appearance

yellow to brownish yellow clarified liquid

Good Manufacturing Practice



Indications

for the treatment of infectious diseases caused by sensitive Gram-positive and Gram-negative bacteria, Rickettsiae, Mycoplasma and other infectious diseases.

Dosage

Intramuscular injection. 0.05-0.1ml per kg body weight of livestock. If necessary, the injection can be repeated at intervals of 48 hours.

Specification

20%



Safety And Packaging

! Cautions

1. Shake well before use
2. Prohibit use in animals with severe liver or kidney dysfunction
3. Prohibit the use on lactating cows.
4. Avoid concurrent use with drugs containing calcium, magnesium, aluminum and iron.
5. Discontinuation period: cattle, sheep, pigs: 28 days, 7 days before weaning.

🔔 Storage

1. Protected from light, sealed and kept in a cool place

📦 Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com