

Lyophilized Powder for Injection *

Amoxicillin Sodium for Injection



Ingredients

Each gram contains: Amoxicillin Sodium 50 mg, with carrier added to make up 1 gram.

Good Manufacturing Practice



- 01 Amoxicillin is primarily used to treat infections caused by Gram-positive and Gram-negative bacteria that are sensitive to penicillin.
- 02 It is applicable for treating symptoms such as fever, anorexia, constipation, recumbency, rapid breathing, and abdominal breathing in poultry and livestock.
- 03 It has shown good efficacy in treating conditions in livestock such as influenza, fever of unknown origin, bacterial dysentery, acute gastroenteritis, swine erysipelas, swine pneumonia, piglet diarrhea, paratyphoid, colibacillosis, brucellosis, mycoplasmosis, leptospirosis, mastitis, endometritis, and agalactia syndrome.

Dosage and Administration

For intramuscular or subcutaneous injection. For livestock, administer 5-10 mg of amoxicillin per kilogram of body weight once daily; or 10-20 mg per kilogram of body weight every other day.

Description

Amoxicillin is a semi-synthetic broad-spectrum penicillin that has antibacterial activity against both Gram-positive and Gram-negative bacteria. Its antimicrobial spectrum includes Campylobacter, Clostridium, Escherichia coli, Erysipelothrix rhusiopathiae, Haemophilus, Pasteurella multocida, Salmonella, penicillin-resistant Staphylococcus, and Streptococcus. The bactericidal action of amoxicillin is achieved by inhibiting bacterial cell wall synthesis. Amoxicillin is primarily excreted through urine, and a portion is also excreted via bile.

Safety And Packaging

Poultry

Escherichia coli disease, Salmonellosis, Fowl cholera, Chicken dysentery, Salpingitis

Contraindications

Animals with severe renal impairment should not use this medication. Concurrent use with tetracyclines, chloramphenicol, macrolides, or lincosamides is contraindicated.

Adverse Reactions

Some livestock may experience allergic reactions, such as edema, although this is relatively rare.



Withdrawal Period

Slaughter: 28 days; Milk: 7 days; Eggs: 7 days.

Packing

1g.2g.4g.8g/bottle
Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com