

Injection



10% Enrofloxacin Injection



Function

Antimicrobial

Ingredients

White or yellowish transparent liquid.

Appearance


100 mg of enrofloxacin per ml.

Good Manufacturing Practice



Indications

For the treatment of livestock and poultry diseases caused by sensitive bacterial and mycoplasma infections, such as Escherichia coli disease, Salmonellosis, Pasteurella, Actinobacillus pleuropneumonia, mastitis, Mycoplasma pneumonia and so on.



Usage and dosage

intramuscular injection. One-time dose.
Pigs, cattle, goats and sheep: 0.025 ml per kg body weight;
Dogs, cats and rabbits: 0.025 – 0.05 ml per kg body weight;
2 times a day for 2-3 days.

Specification

10%, 1ml: 0.1g.

Safety And Packaging

Side effects

This product has very low toxicity.

The main side effects include

1. may affect the normal development of bones in young animals, resulting in pain or lameness.
2. May cause vomiting, loss of appetite and diarrhea.
3. may develop erythema, itching, hives or photosensitivity reactions.
4. May occasionally cause allergic or seizure reactions in cats and dogs.

Rest period

Cattle, rabbits and sheep: 14 days; pigs: 10 days.



Storage

Protect from light and keep sealed.

Packing

100ml/bottle

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613
E-mail: hebeijinfeng@gmail.com