



Fenbendazole Tablets



Good Manufacturing Practice



Fenbendazole is a broad-spectrum antiparasitic drug used primarily for the treatment of parasitic infections in a wide range of animals, including:

- Nematodes: e.g. roundworms, hookworms, whipworms, etc.
- Cestodes: e.g. Moniezia tapeworms, etc.
- Trematodes: e.g. liver flake trematode, etc.
- · Larvae and eggs: with the effect of killing eggs.
- Suitable for many kinds of animals such as pigs, cattle, sheep, horses, dogs, cats,
- poultry and so on.



Usage and Dosage

For internal use: Dosage varies according to the weight and type of parasite in different animals.

Pig: 5mg per kg body weight, use for 3 days. Cattle, sheep: 0.1-0.15g per kilogram of body weight for 3 days. Dogs, cats: 25-50mg per kilogram of body weight for 3 days. Poultry: 10-50mg per kg body weight.

Safety And Packaging

Opening Dormancy period

- 1. Single dose ineffective: For animals such as dogs and cats, 3 consecutive days of use is usually required.
- 2. Prohibitions Prohibited in horses intended for food. Not to be used in lactating cows and sheep. Prohibited during the first 45 days of
- 3. POSSIBLE SIDE EFFECTS: Fenbendazole does not usually produce adverse reactions at regular doses, but high doses may cause allergic reactions or vomiting.

Storage conditions: Shielded from light, airtight and kept in a dry place.



Dormant period

Cattle, sheep: 21 days. Pig: 3 days. Poultry: 28 days. Milk abandonment period: 7 days. Egg abandonment period: 7 days.



Shelf life

2 years



Packing

5 tablets/plate

Packaging specifications can be adjusted according to customer requirements.

Feel Free to Contact Us for Your Needs!

Phone/WhatsApp: +86 15321587613 E-mail: hebeijinfeng@gmail.com